

# A Multicanonical Algorithm and the Surface Free Energy in SU(3) Pure Gauge Theory

B. Grossmann<sup>1</sup>, M. L. Laursen<sup>1</sup>,  
T. Trappenberg<sup>1,2</sup>, U.-J. Wiese<sup>3,a</sup>

<sup>1</sup>HLRZ,c/o Kfa Juelich, D-5170 Juelich, Germany

<sup>2</sup>Inst. f. Theor. Physik, RWTH Aachen, D-5100 Aachen, Germany

<sup>3</sup>Inst. f. Theor. Physik, Univ. Bern, CH-3012 Bern, Switzerland.

## Abstract

We present a multicanonical algorithm for the SU(3) pure gauge theory at the deconfinement phase transition. We measure the tunneling times for lattices of size  $L^3 \times 2$  for  $L = 8, 10$ , and  $12$ . In contrast to the canonical algorithm the tunneling time increases only moderately with  $L$ . Finally, we determine the interfacial free energy applying the multicanonical algorithm.

## 1 Introduction

For the analysis of systems with first order phase transitions—as e.g. quenched QCD at the deconfinement temperature—surface effects are of crucial importance. They determine the physical dynamics and lead to supercritical slowing down in computer simulations at the critical point. Configurations which include an interface are exponentially suppressed by their free energy which results in an exponential increase of the tunneling time. Recently the multicanonical algorithm was proposed in order to overcome this supercritical slowing down [1]. Here we present a variant of this method for the SU(3) gauge theory with the Wilson action. We have used lattices of size  $L^3 \times L_T$  with  $L_T = 2$  and  $L = 8, 10$ , and  $12$  for which there is a strong first order phase transition. We achieve a considerable reduction in the tunneling time with only a modest increase in computer time. We apply this algorithm to the determination of the interfacial free energy

$$F_{cd} = \alpha_{cd} A \quad (1.1)$$

of a confinement-deconfinement interface of area  $A$ ,  $\alpha_{cd}$  being its surface tension. For this we made use of Binder's histogram method which in contrast to the methods used in [2, 3]

---

<sup>a</sup>Supported by Schweizer Nationalfond

generates interfaces dynamically rather than by introducing artificial external parameters. In that respect, this approach is similar to transfer matrix methods [4].

## 2 Monte Carlo Simulations at a First Order Phase Transition

As mentioned above, MC simulations close to first order phase transitions suffer from supercritical slowing down, i.e. the tunneling time  $\tau_L$  for a system of volume  $V = L^3$  is expected to be proportional to the inverse of the probability of a system with interfaces (actually there will be two interfaces because of the boundary conditions),

$$\tau_L \propto \exp(2L^2 \frac{\alpha}{T}) \quad (2.1)$$

and therefore diverges exponentially with the area  $A = L^2$  of an interface.

In order to overcome this problem, the multicanonical algorithm does not sample the configurations with the canonical Boltzmann weight

$$\mathcal{P}_L^{can}(S) \propto \exp(\beta S), \quad (2.2)$$

where  $S = 1/3 \sum_{\square} tr U_{\square}$  is the Wilson action in four dimensions, but rather with a modified weight

$$\mathcal{P}_L^{mc}(S) \propto \exp(\beta_L(S)S + \alpha_L(S)). \quad (2.3)$$

The coefficients  $\alpha_L$  and  $\beta_L$  are chosen such that the probability  $P_L$  (not to be confused with the Boltzmann weights) is increased for all values of the action in between the two maxima  $S_L^{max,1}$  and  $S_L^{max,2}$ , as shown schematically in figure 1. Finally data are analyzed from the multicanonical samples by reweighting with  $\exp((\beta - \beta_L(S))S - \alpha_L(S))$ .

In order to approximate the weights  $\mathcal{P}_L^{mc}$  leading to the distribution  $P_L^{mc}$  of figure 1, we start from some good estimate of the canonical distribution  $P_L$  (see below) at the coupling  $\beta_L^c$  which corresponds to equal weight in both phases. This will lead to different heights of the two peaks. Then we take a partition

$$\begin{aligned} S_L^0 = -6L^3 L_t &< S_L^1 \equiv S_L^{max,1} < S_L^2 < \dots < S_L^{N/2} \equiv S_L^{min} < \dots \\ \dots &< S_L^{N-1} \equiv S_L^{max,2} < S_L^N = 6L^3 L_t \end{aligned} \quad (2.4)$$

of the interval  $-6L^3 L_t \leq S \leq 6L^3 L_t$ . The coefficients  $\alpha_L$  and  $\beta_L$  are chosen to be constants  $\alpha_L^k$  and  $\beta_L^k$  in the intervals  $[S_L^k, S_L^{k+1})$  such that  $\ln P_L^{mc}$  interpolates the linear function

$$\ln P_L^{max,1} + (\ln P_L^{max,2} - \ln P_L^{max,1}) \cdot \frac{S - S_L^{max,1}}{S_L^{max,2} - S_L^{max,1}}. \quad (2.5)$$

continuously between the points  $S_L^k$  and  $S_L^{k+1}$  for  $k = 1$  to  $N - 2$ . The Boltzmann weight is identical to the canonical one in the first and last interval. We arrive at

$$\beta_L^k - \beta = \begin{cases} 0 & , k = 0 \\ \delta\beta + \ln\left(\frac{P_L(S_L^k)}{P_L(S_L^{k+1})}\right) / (S_L^{k+1} - S_L^k) & , k = 1, \dots, N - 2 \\ 0 & , k = N - 1 \end{cases} \quad (2.6)$$

where  $\delta\beta \equiv \ln \left( \frac{P_L^{max,2}}{P_L^{max,1}} \right) / (S_L^{max,2} - S_L^{max,1})$ , and the  $\alpha_L^k$  are given by

$$\alpha_L^{k+1} \equiv \alpha_L^k + (\beta_L^k - \beta_L^{k+1}) S_L^{k+1}, \quad \alpha_L^0 = \delta\beta \cdot S_L^{max,1}. \quad (2.7)$$

The partition  $\{S_L^k\}$  which is in principle arbitrary is defined by demanding

$$\frac{P_L(S_L^{k+1})}{P_L(S_L^k)} = \begin{cases} 1/r_1 & , k = 1, \dots, N/2 - 1 \\ r_2 & , k = N/2, \dots, N - 2, \end{cases} \quad (2.8)$$

such that  $r_1^{N/2-1} = P_L^{max,1}/P_L^{min}$  and  $r_2^{N/2-1} = P_L^{max,2}/P_L^{min}$ . This generalizes the formulas given in [1] to the case  $P_L^{max,1} \neq P_L^{max,2}$  by introducing a  $\delta\beta \neq 0$ . In addition the specific choice for  $r_1$  and  $r_2$  assures that the probability of  $S_L^{min}$  is lifted by the correct amount.

We apply this algorithm to the  $SU(3)$  pure gauge theory at the deconfinement phase transition. The multicanonical data sampling was done with a 5-hit Metropolis algorithm as well as with a Creutz heat bath algorithm modified according to equ. (2.3). In both cases three independent  $SU(2)$ -subgroups are updated following the idea of ref. [5]. Because of the dependence of  $\alpha_L$  and  $\beta_L$  on the total action  $S$ , the update of the active link has to be done in scalar mode. However, most of the computation time is needed for the calculation of the staples surrounding the active link, and this is still vectorizable. In addition, for the Creutz algorithm we profited from the fact that as long as the action stays within one interval  $[S_L^k, S_L^{k+1})$  there is no dependence of  $\alpha_L$  and  $\beta_L$  on  $S$ . Therefore, if the intervals are not too small even the update can be vectorized partially, since one hardly ever crosses an interval boundary.

In table 2 the update times for the various algorithms for one processor of a CRAY-YMP are given for a  $10^3 \times 2$  lattice using  $N = 14$  intervals for  $\mathcal{P}_L^{mc}$ . As can be seen the CPU time per link and update is increased by only about 50%. In our simulations we use

Table 1. CPU time per link update for canonical and multicanonical versions of a Metropolis and a Creutz heatbath algorithm.

| Algorithmus        | Metr. | Metr.(mc) | Creutz | Creutz(mc) |
|--------------------|-------|-----------|--------|------------|
| $t_{upd}(\mu sec)$ | 37    | 53        | 31     | 52         |

$C$ -periodic boundary conditions in the three spatial directions (see [6]). This suppresses the two deconfined phases with nonvanishing expectation value of the imaginary part of the Polyakov line. Then only the confined and one deconfined phase coexist at the critical temperature  $T_c$ . For finite volumes we define  $T_L^c$  (or  $\beta_L^c$  resp.) by demanding equal weight for these two phases [7], i.e.

$$P_L(S \leq S_L^{min}) \equiv P_L(S \geq S_L^{min}). \quad (2.8)$$

We typically do 50000 updates for  $\beta \approx \beta_L^c$ . One sweep consists of one Metropolis (resp. Creutz) update step plus four overrelaxation steps. For the smaller lattices  $L = 8$  and 10 the parameters for the multicanonical algorithm were taken from canonical simulations. For

$L = 12$  where the canonical algorithm is no longer efficient due to the increasing tunneling time, we took the parameters from an appropriate finite size scaling extrapolation of the distributions for the  $10^3 \times 2$  lattice. In all cases we take  $N = 14$  intervals.

Figure 2 shows part of the history of the average Polyakov line as a function of the number of sweeps for the various algorithms. From the full histories we extract the tunneling times  $\tau_L$ , defined as one fourth of the number of sweeps needed for a full tunneling cycle. The tunneling times  $\tau_L$  in table 2 indicate that the increase with  $L$  for the standard algorithm is effectively reduced. Already for  $L = 10$  the gain in  $\tau_L$  is much larger than the overhead in CPU time per update which is due to scalar operations.

Table 2. Tunneling times  $\tau_L$  for the various algorithms and lattices of size  $L^3 \times 2$ .

| L  | $\beta$ | Metr. | Metr.(mc) | Creutz(mc) |
|----|---------|-------|-----------|------------|
| 8  | 5.094   | 300   | 200       | –          |
| 10 | 5.095   | 1500  | 400       | 400        |
| 12 | 5.0928  | –     | 700       | 700        |

lattices of

Thus the multicanonical algorithm provides an efficient tool for Monte Carlo studies of the deconfinement phase transition. As an application, we present in the next section the determination of the interfacial free energy  $F_S$ .

### 3 The Interfacial Free Energy

The configurations with  $S = S_L^{min}$  are expected to contain two parallel interfaces of area  $L^2$  between the confined and the deconfined phase. Therefore they should be suppressed by the interfacial free energy  $\alpha L^2$ . We expect additional power law corrections due to capillary wave fluctuations and translational Goldstone modes of the interfaces, such that

$$P_L^{min} \propto L^x \exp\left(-2L^2 \frac{\alpha}{T}\right) \quad (3.1)$$

(see also ref.[8]). The probability distribution should be flat around  $S_L^{min}$  for non-interacting surfaces due to translations of the interfaces relative to each other.

We extract the probability distribution  $P_L$  for the plaquette and the real part of the Polyakov line from our multicanonical simulations for  $L = 8, 10$ , and  $12$  (see figure 3). They agree reasonably with those obtained in ref. [9] for periodic boundary conditions. However, there is no plateau for the probabilities around  $S_L^{min}$ . This is likely to be due to interfacial interactions which will be reduced by using a larger extension in one of the spatial directions. This is left for further investigations.

Nevertheless, we consider the two quantities

$$F_L^{(1)} \equiv \frac{1}{2L^2} \ln \frac{\overline{P_L^{max}}}{P_L^{min}} \quad (3.2)$$

and

$$F_L^{(2)} \equiv -\frac{1}{2L^2} \ln P_L^{min} \quad (3.3)$$

where  $\overline{P}_L^{max} \equiv \frac{1}{2}(P_L^{max,1} + P_L^{max,2})$ . They should converge to  $\frac{\alpha}{T}$  for  $L \rightarrow \infty$  for non-interacting surfaces. We plot them as functions of  $1/L^2$  in figure 4. We find a reasonable agreement between the values taken from the plaquette and Polyakov line distributions. As expected, the discrepancy between  $F_L^{(1)}$  and  $F_L^{(2)}$  decreases for increasing  $L$ . Still, compared to the values of  $\alpha$  obtained by different methods [2, 4] the extrapolated result of our data is too small. Simulations on asymmetric lattices should clarify this point.

In conclusion we have demonstrated the power of the multicanonical algorithm for SU(3) pure gauge theory. The tunneling time  $\tau_L$  is reduced considerably while the update time is only increased by less than 50% compared to standard Metropolis or heat bath algorithms. This reduces the computer time needed for investigations of the deconfinement phase transition like the one presented here for the interfacial free energy. Using cylindrical lattices will probably reduce the interactions between the interfaces thereby allowing the extraction of the interface tension.

## References

- [1] B. A. Berg, U. Hansmann und T. Neuhaus, *preprint FSU-SCRI-91-125*;  
B. A. Berg, T. Neuhaus, *preprint BI-TP 91/08*; *Phys. Lett. B267* (1991) 249-253;  
*Phys. Rev. Lett.* 68 (1992) 9.
- [2] S. Huang, J. Potvin, C. Rebbi und S. Sanielevici, *Phys. Rev. D42* (1990) 2864.  
R. Brower, S. Huang, J. Potvin, C. Rebbi und J. Ross, *preprint BUHEP-91-22*.  
R. Brower, S. Huang, J. Potvin, und C. Rebbi, *preprint BUHEP-92-3*.
- [3] K. Kajantie, L. Kärkkäinen und K. Rummukainen, *Nucl. Phys. B333* (1990) 100.
- [4] B. Großmann, M. L. Laursen, T. Trappenberg and U.-J. Wiese, *HLRZ-preprint* in preparation.
- [5] N. Cabibbo and E. Marinari, *Phys. Lett.* 119B (1982), 387.
- [6] A. S. Kronfeld, U. J. Wiese, *Nucl. Phys. B357* (1991) 521;  
U.-J. Wiese, *Nucl. Phys. B375* (1992) 45.
- [7] C. Borgs and W. Janke, *Phys. Rev. Lett.* 68 (1992) 1738.
- [8] K. Binder, *Z. Phys. B43* (1981) 119 and *Phys. Rev. A25* (1982) 1699.
- [9] W. Janke, B. A. Berg und M. Katoot, *preprint FSU-SCRI-92-40*.

## 4 Figure Captions

Figure 1 Schematic representation of the probability distributions  $P_L$  (straight line) and  $P_L^{mc}$  (dashed line) corresponding to the canonical Boltzmann weight  $\mathcal{P}_L^{can}$  and the multicanonical one  $\mathcal{P}_L^{mc}$ . For  $S < S_L^{max,1}$  and  $S > S_L^{max,2}$ ,  $P_L$  and  $P_L^{mc}$  (normalized to their maximum value) are identical.

Figure 2 Evolution of the average Polyakov line as a function of the total number of sweeps for a  $10^3 \times 2$  lattice for (A) a canonical Metropolis algorithm, (B) a multicanonical Metropolis algorithm, and (C) a multicanonical Creutz algorithm.

Figure 3 Distribution for the average plaquette and the real part of the Polyakov line for lattice sizes  $L = 8, 10$  and  $12$ . from the average versus  $\ln L/L^2$ . indicated by a diamond,

Figure 4 The free energies  $F_L^{(1)}$  and  $F_L^{(2)}$  from the average plaquette and the real part of the Polyakov line versus  $1/L^2$ . The value for  $\alpha/T$  from reference [2] is indicated by a diamond, the value from reference [3] by a filled square, and the value from reference [4] by a triangle.